

Anatomy and Physiology

Nervous System

www.physiotherapyphd.com

The Cerebellum

The cerebellum, which stands for “little brain”, is a structure of the central nervous system. It has an important role in motor control, with cerebellar dysfunction often presenting with motor signs. In particular, it is active in the coordination, precision and timing of movements, as well as in motor learning.

During embryonic development, the anterior portion of the neural tube forms three parts that give rise to the brain and associated structures:

- Forebrain (prosencephalon)
- Midbrain (mesencephalon)
- Hindbrain (rhombencephalon)

The hindbrain subsequently divides into the **metencephalon** (superior) and the **myelencephalon** (inferior). The cerebellum develops from the metencephalon division.

This article will focus on the anatomy of the cerebellum. It will provide a brief overview of its functions and development, and finally it will highlight the clinical relevance of cerebellar disorders.

Location

The cerebellum is located at the back of the brain, immediately inferior to the **occipital** and **temporal lobes**, and within the posterior cranial fossa. It is separated from these lobes by the **tentorium cerebelli**, a tough layer of dura mater.

It lies at the same level of and posterior to the **pons**, from which it is separated by the fourth ventricle.

Anatomical structure and divisions

The cerebellum consists of two hemispheres which are connected by the **vermis**, a narrow midline area. Like other structures in the central nervous system, the cerebellum consists of grey matter and white matter:

- **Grey matter** – located on the surface of the cerebellum. It is tightly folded, forming the cerebellar cortex.
- **White matter** – located underneath the cerebellar cortex. Embedded in the white matter are the four cerebellar nuclei (the dentate, emboliform, globose, and fastigi nuclei).

Anatomical Lobes

There are three anatomical lobes that can be distinguished in the cerebellum; the anterior lobe, the posterior lobe and the flocculonodular lobe. These lobes are divided by two fissures – the **primary fissure** and **posterolateral fissure**.

Zones

There are three cerebellar zones. In the midline of the cerebellum is the **vermis**. Either side of the vermis is the **intermediate zone**. Lateral to the intermediate zone are the **lateral hemispheres**. There is no difference in gross structure between the lateral hemispheres and intermediate zones

Functional Divisions

The cerebellum can also be divided by function. There are three functional areas of the cerebellum – the cerebrocerebellum, the spinocerebellum and the vestibulocerebellum.

- **Cerebrocerebellum** – the largest division, formed by the lateral hemispheres. It is involved in planning movements and motor learning. It receives inputs from the cerebral cortex and pontine nuclei, and sends outputs to the thalamus and red nucleus. This area also regulates coordination of muscle activation and is important in visually guided movements.
- **Spinocerebellum** – comprised of the vermis and intermediate zone of the cerebellar hemispheres. It is involved in regulating body movements by allowing for error correction. It also receives proprioceptive information.
- **Vestibulocerebellum** – the functional equivalent to the flocculonodular lobe. It is involved in controlling balance and ocular reflexes, mainly fixation on a target. It receives inputs from the vestibular system, and sends outputs back to the vestibular nuclei

The cerebellum receives its blood supply from three paired arteries:

- Superior cerebellar artery (SCA)
- Anterior inferior cerebellar artery (AICA)
- Posterior inferior cerebellar artery (PICA)

For more notes click on:
www.physiotherapyphd.com